CERN-PH-TH-2015-080 — ZU-TH 09/15 — MCnet-15-08

Hadronic Higgs production through NLO+PS in the SM, the 2HDM and the MSSM

Hendrik Mantler[a], Marius Wiesemann[b]

[a] *TH Division, Physics Department, CERN, CH-1211 Geneva 23, Switzerland*

[b] *Physik-Institut, Universität Zürich, 8057 Zürich, Switzerland*

hmantler@cern.ch
mariusw@physik.uzh.ch

arXiv:1504.06625v1 [hep-ph] 24 Apr 2015

Abstract

The NLO cross section of the gluon fusion process is matched to parton showers in the MC@NLO approach. We work in the framework of `MadGraph5_aMC@NLO` and document the inclusion of the full quark-mass dependence in the SM as well as the state-of-the-art squark and gluino effects within the MSSM embodied in the program `SusHi`. The combination of the two programs is realized by a script which is publicly available and whose usage is detailed. We discuss the input cards and the relevant parameter switches. One of our focuses is on the shower scale which is specifically important for gluon-induced Higgs production, particularly in models with enhanced Higgs-bottom Yukawa coupling.

1 Introduction

Higgs production proceeds predominantly through gluon fusion in a large number of theories, including the Standard Model (SM). The recently discovered resonance [1, 2] in searches for a Higgs boson is fully consistent with the SM picture[1], so far. Still, the measured Higgs boson may be embedded in an enlarged Higgs sector with respect to the one of the SM which predicts only a single physical particle breaking the electro-weak symmetry. Two-Higgs Doublet Models (2HDM's) such as the Minimal Supersymmetric SM (MSSM) are among the most popular theories with enlarged Higgs sectors. Such theories inevitably require the existence of further physical Higgs particles. A 2HDM predicts three neutral Higgs bosons: a light (h) and a heavy (H) scalar, and a pseudo-scalar (A); as well as two charged Higgs particles ($H^{\pm}$). Almost all 2HDM and MSSM scenarios that are in agreement with the experimental bounds feature a light scalar which is SM-like in its couplings to vector bosons and fermions, while the other Higgs bosons are heavier and, therefore, escaped detection up to now. Indeed, the experimental search for other Higgs resonances is one of the major focuses regarding the discovery of physics beyond the SM (BSM) in the second run of the Large Hadron Collider (LHC).

Higgs production through gluon fusion is mediated by a colored particle. In the SM, the top quark gives the dominant contribution to the cross section [6–8]. While also the bottom quark gives a sizable contribution, the effects due to other quarks are small and therefore usually neglected. In the 2HDM and the MSSM the Higgs-bottom Yukawa coupling can be enhanced with respect to the one of the top quark and the bottom loop may even constitute the dominant contribution to the cross section. In those models it is stringently required to include the bottom-quark contribution.[2] The gluon fusion cross section is known at the next-to-leading order (NLO) in the SM including top- and bottom-mass effects [18,19], in the 2HDM and in the MSSM including contributions from squarks and gluinos [20–34]. For the top quark, an effective field theory approach can be applied in which the top quark is considered to be infinitely heavy and can be integrated out from the full theory. In this approximation, Higgs production has been calculated up to the next-to-NLO (NNLO) inclusively [35–37] as well as fully-differential [38–40]. Electro-weak contributions and effects beyond NNLO in the heavy-top approximation have been studied in Refs. [41–50] for example, while there was a large effort [51, 52] to push the accuracy to next-to-NNLO (N^{3}LO) which has been succeeded very recently [53]. Finite top-mass effects have been shown to be small for both the inclusive cross section at the NNLO [54–59] and differential quantities [60, 61] as long as no kinematical scale (such as the transverse momentum of a particle) that is not integrated out exceeds the top-mass threshold.

The full dependence of the top- and the bottom-mass at the NLO has been included so far in a POWHEG-type [62] matching to parton showers (PSs) [33], the analytically resummed transverse momentum spectrum of the Higgs boson at NLO+NLL [63], a MC@NLO-type [64]

[1] See Refs. [3–5] for a theoretical overview.

[2] Note that in theories with an enhanced bottom Yukawa the associated Higgs production with bottom quarks becomes relevant, see Refs. [9–17] and references therein.

matching to the Herwig Monte Carlos [65–67], the NNLO+NNLL jet-vetoed [68] and the fully-differential NNLO [69] cross section; and in some approximated form recently also in the NNLOPS approach [70, 71]. Furthermore, the 2HDM as well as supersymmetric effects from squarks and gluinos within the MSSM [20–34] have been implemented in the first two approaches from that list [33,72]. In this manuscript, we report on a new implementation of NLO QCD corrections in the SM, 2HDM and MSSM applying the MC@NLO-type matching to both Herwig and Pythia showers. We work in the framework of MadGraph5_aMC@NLO [73] and combine its capabilities with the corresponding amplitudes provided by SusHi [74]. The linking of SusHi to MadGraph5_aMC@NLO is realized by a script[3] aMCSusHi. Its usage as well as the application of the combined code to obtain cross section predictions in the SM, the 2HDM and the MSSM is detailed in this paper.[4]

The manuscript is organized as follows: In Section 2 we present a brief overview of the elements of the computation at hand. Section 3 is dedicated to introduce aMCSusHi and is separated in three parts which cover: how to use the script (Section 3.1), how to run the resulting code (Section 3.2) and how to treat the shower scale (Section 3.3). We will show a brief application of the code to phenomenological results in Section 4 and conclude in Section 5.

2 Outline of the calculation

The goal of this paper is to present a tool which allows for the computation of arbitrary infra-red safe differential observables at both the parton- and the hadron-level for the production of neutral Higgs bosons via gluon fusion in the SM, the 2HDM and the MSSM by matching the NLO cross section to a shower.

The relevant NLO matrix elements are taken from Ref. [74], which include both SM-like contributions and sbottom, stop and gluino effects. Examples of corresponding Feynman diagrams are illustrated in Fig. 1. They are combined and matched to a parton shower by the well-known MC@NLO-method. The matched cross section in the MC@NLO framework can be written symbolically as:

$$\begin{aligned}\left(\frac{d\sigma}{dO}\right)_{\text{MC@NLO}} = &\int d\Phi_n \left[B_n + V_n + \int d\Phi_1^{\text{MC}}\, K_{n+1}^{\text{MC}}\right] \mathcal{I}_n^{\text{MC}}(O) \\ &+ \int \left[d\Phi_{n+1} R_{n+1} - d\Phi_{n+1}^{\text{MC}} K_{n+1}^{\text{MC}}\right] \mathcal{I}_{n+1}^{\text{MC}}(O)\,,\end{aligned} \tag{1}$$

where B_n determines the Born-level cross section, V_n the virtual (including mass factorization) and R_{n+1} the real corrections; K_{n+1}^{MC} is the Monte Carlo subtraction term, with the same IR poles as R_{n+1}, the Monte Carlo phase space $d\Phi_{n+1}^{\text{MC}}$ tends to $d\Phi_{n+1}$ in the IR limits, and $d\Phi_1^{\text{MC}} = d\Phi_{n+1}^{\text{MC}}/d\Phi_n$. The quantity $\mathcal{I}_n^{\text{MC}}(O)$ is the shower spectrum for observable O, as obtained by running the shower starting from an n-body configuration.

[3] aMCSusHi can be downloaded under https://cp3.irmp.ucl.ac.be/projects/madgraph/wiki/HSushi#no1.

[4] SusHi has recently been extended to include NMSSM Higgs production [75], which may be made available in aMCSusHi in the future.



Figure 1: A sample of Feynman diagrams for $gg \to \phi$ contributing to the NLO cross section; (a-c) LO, (d-g) virtual and (h-i) real corrections. The graphical notation for the lines is: solid straight $\widehat{=}$ quark; spiraled $\widehat{=}$ gluon; dashed $\widehat{=}$ scalar (squark or Higgs); spiraled with line $\widehat{=}$ gluino.

The cross section at Born-level is derived from the LO diagrams for $gg \to \phi$ where $\phi \in \{h, H, A\}$, see e.g. Fig. 1 (a)-(c). The NLO virtual and real corrections are governed by diagrams like the ones shown in Fig. 1 (d)-(g) and Fig. 1 (h)-(i), respectively, and similar ones with quark loops replaced by squark loops.

Eq. (1) is implemented for all standard parton showers [76–81] in the fully-automated framework `MadGraph5_aMC@NLO`. This code determines NLO QCD corrections to arbitrary scattering processes at the LHC. On the basis of UFO models [82], the code even allows to carry out computations in any theory beyond the SM in a general manner as soon as the renormalization is known and implemented in a UFO model, see Refs. [73, 83–85] for further information. However, the Higgs production mode through gluon fusion is a special one being loop-induced already at the LO. Such processes cannot be handled in a fully-automated manner by any code to date, since it requires the automation of two-loop amplitudes which is beyond current technology. Therefore, we have treated Higgs production through gluon fusion in the SM, 2HDM and MSSM as a special case, by linking the relevant amplitudes from `SusHi`. Furthermore, as far as the MSSM is concerned

SusHi requires a link to FeynHiggs [86–96] which evaluates the corresponding Higgs masses and couplings in user defined scenarios. Setting up the SusHi amplitudes in MadGraph5_aMC@NLO is handled by a publicly available script called aMCSusHi, which is automated to create the $gg \to \phi$ process folder; download SusHi and FeynHiggs; compile, install and link them; and replace the relevant amplitudes in the process folder. In the upcoming section, we describe the application of the script and explain the necessary steps to obtain phenomenological results.

3 aMCSusHi script

This code is based on MadGraph5_aMC@NLO and SusHi. For further information on theses codes we refer the reader to corresponding publications [73, 74]. After using the script to set up the code, we will focus on the relevant user inputs to obtain phenomenological predictions for Higgs cross sections in the SM, the 2HDM and the MSSM.

3.1 Usage of the script

The aMCSusHi script is available for download from the website https://cp3.irmp.ucl.ac.be/projects/madgraph/wiki/HSushi#no1. It is fully automatic in setting up the $gg \to \phi$ process folder which includes downloading, installing and linking FeynHiggs and SusHi. At first, a dummy HEFT process folder for $gg \to h$ in the five-flavor scheme at NLO (without the virtuals) is created;[5] then, the HEFT amplitudes in MadGraph5_aMC@NLO are consistently replaced by the ones from SusHi (including the virtuals). The set-up requires only a single call of the aMCSusHi script:

```
> ./set_up_ggH_MSSM_script.pl <ggH-folder> [<FeynHiggs-folder> [<SusHi-folder>]]
```

The first argument is mandatory and determines the path to the process folder for $gg \to \phi$ that is generated by the script. The only requirement is that this folder has to be defined as a sub-folder of the MadGraph5_aMC@NLO directory. The second and third arguments are optional. If there are compiled versions of FeynHiggs and SusHi available on your computer you can give the names of the folders that contain the files libFH.a and libsushi.a, respectively. When executed with two (one) arguments the script will ask whether it should automatically download and install SusHi (and FeynHiggs). The script will always download the latest versions of these codes. While running, the script requires some user inputs: It asks whether or not SusHi (and FeynHiggs) should be downloaded, in which folder they should be installed (default is inside the <ggH-folder>) or where to find SusHi (and FeynHiggs) if already installed. The user is simply required to follow these on-screen instructions. Furthermore, the script creates log-files in the working directory for the download ("XX_curl.log"), the configure command ("XX_conf.log") and the compilation ("XX_make.log"), where XX=FH for FeynHiggs and XX=SusHi for SusHi.

[5] See the README file how this can be easily obtained with the MadGraph5_aMC@NLO script.

These files are supposed to give complementary information for any kind of troubleshooting.[6] Further information about the `aMCSusHi` script provides the README file.

3.2 Running the code

Once the $gg \to \phi$ process folder has been set up by the script, the run can be started directly from the <ggH-folder> by typing

```
> ./bin/generate_events
```

and following the usual steps in `MadGraph5_aMC@NLO` to choose the run-modes (order, shower or fixed-order, madspin). Before running the code one may want to modify the input settings. In the following we will discuss the differences between `aMCSusHi` and the ordinary `MadGraph5_aMC@NLO` code regarding the input files.

The input cards can be found under <ggH-folder>/Cards/, where the param_card.dat, run_card.dat and shower_card.dat contain all the essential information. The run card controls the usual parameters, e.g. the renormalization ($\mu_{\rm R}$) and factorization scale ($\mu_{\rm F}$). Note, that by default the flags `fixed_ren_scale` and `fixed_fac_scale` are set to false, so that these two scales are chosen on an event-wise basis. Their values are specified in the last routine of <ggH-folder>/SubProcesses/setscales.f which, due to the default option `dynamical_scale_choice` $= -1$ in the run card, sets $\mu_{\rm R} = \mu_{\rm F} = H_T/2 \equiv 1/2 \sum_i (m_i^2 + p_T^2(i))^{1/2}$, where i runs over all final state particles and m_i and $p_T(i)$ are their mass and transverse momentum, respectively. This choice is reasonable, since it respects effects from hard radiation and corresponds to a value of $m_\phi/2$ in the soft/collinear limit which is the current recommendation for the total inclusive $gg \to \phi$ cross section [3, 5].

Also the shower card in `aMCSusHi` contains no new information and has the usual functionality. Since `MadGraph5_aMC@NLO` supports all standard parton showers, for the first time `Pythia6` and `Pythia8` can be applied at NLO+PS to SM Higgs production in the full theory in the MC@NLO framework. In general, it is advisable to apply the most recent versions of the showers for meaningful physics runs.

The param_card.dat, on the other hand, receives some significant changes by the `aMCSusHi` script. The new version basically combines the orignial parameter card from `MadGraph5_aMC@NLO` with the input file from `SusHi` which are both written in the SUSY Les Houches accord (SLHA) format [97] and, therefore, easily connectable. We will address the different options in the param_card.dat in more detail, since there are a number of changes and some of the original parameters loose their functionality. In the SLHA format inputs are organized in blocks which have different entries that are characterized by a number. For simplicity, we introduce the following short-hand notation: `Block example[i]` corresponds to entry i in `Block example`.

[6] Please note that even in cases where the compilation of `FeynHiggs` or `SusHi` fails, the linking may still work fine as long as the files libFH.a and libsushi.a have been created in the corresponding library folders.

E. g., entry 25 of `Block mass` (`Block mass[25]`) in the SLHA format is devoted to the Higgs mass in the SM, which is required as an input in the param_card.dat. A typical parameter card of `aMCSusHi` in the SM is shown below:

```
###################################
## INFORMATION FOR MASS
###################################
Block mass
   15 1.777000e+00 # MTA
   23 9.118800e+01 # MZ
   25 1.250000e+02 # MH -- only effective if FEYNHIGGS Block is absent
[...]

###################################
## INFORMATION FOR SMINPUTS
###################################
Block sminputs
    1 1.325070e+02 # aEWM1
    2 1.166390e-05 # Gf
    3 1.180000e-01 # aS
# additional information needed for SusHi
    4 9.118760e+01 # m_Z(pole)
    5 0.416000e+01 # m_b(m_b)  -- only used if m_b is not on-shell
    6 1.730000e+02 # m_t(pole) -- top mass is set here

###################################
## INFORMATION FOR YUKAWA
###################################
Block yukawa
   15 1.777000e+00 # ymtau

###################################
## INFORMATION FOR DECAY
###################################
DECAY   6 1.491500e+00 # WT
DECAY  23 2.441404e+00 # WZ
DECAY  24 2.047600e+00 # WW
DECAY  25 6.382339e-03 # WH
DECAY 9000006 6.382339e-03 # WH1
[...]

###################################
## INFORMATION FOR SUSHI
###################################
Block sushi
  1   0          # model: 0 = SM, 1 = MSSM, 2 = 2HDM
  2   0          # 0 = light Higgs (h), 1 = pseudoscalar (A), 2 = heavy Higgs (H)
Block renormbot # Renormalization of the bottom sector
  1   0          # m_b used for bottom Yukawa:  0 = OS, 1 = MSbar(m_b), 2 = MSbar(muR)
  4   4.75d0     # mbOS fixed -- used if m_b is on-shell (default)
Block factors
  1   0.d0       # factor for yukawa-couplings: c
  2   1.d0       # t
  3   1.d0       # b
```

Most of the inputs are self-explanatory due to the comments initiated by the hash symbol **#** after the entries. Furthermore, the standard SLHA blocks match the universal convention of Ref. [97].

Some of the inputs, though, require further comments. In the `Block mass` all parameters have the expected function, except for the top and the bottom mass, `Block mass[6]` and `Block mass[5]`, respectively. While the former only affects and is required for the shower, the latter can be omitted completely. Instead, due to the link to `SusHi` the top mass that is used for the top loop and the top Yukawa is set in `Block sminputs[6]` and is expected to be on-shell. For the bottom mass, on the other hand, `SusHi` allows for three different choices: on-shell scheme or $\overline{\text{MS}}$ scheme with $m_b(m_b)$ or $m_b(\mu_\text{R})$, which can be switched in `Block renormbot[1]` by a value between 1 and 3.[7] Also here the recommendation is to use the on-shell scheme which according to Refs. [18, 98] assures the cancellation of large logarithms $\ln(m_h/m_b)$ at NLO QCD, while the $\overline{\text{MS}}$ scheme does not, due to an incomplete resummation of these terms. The on-shell m_b value is determined by `Block renormbot[4]`, while when the $\overline{\text{MS}}$ scheme is chosen the corresponding input of $m_b(m_b)$ is set in `Block sminputs[5]`. The other entries of `Block sminputs` again have the same impact as in the usual `MadGraph5_aMC@NLO` code. The same is true for `Block yukawa`. For the decay of light and heavy Higgs bosons one may specify a finite width of the Higgs boson in the respective BSM scenario by using `Decay 25` irrespective of wether the light or the heavy Higgs boson is considered. At this point we shall remark that the particle identification number (PID) in the generated event files is always `25` regardless of the Higgs boson under consideration.[8] This is irrelevant for the production (which is correctly computed through the `SusHi` amplitudes), but plays a role for the decay where the shower will consider particle `25` to be the light Higgs, which is indeed fine for any scalar Higgs, but a problem for pseudo-scalar ones. Therefore, decays of a pseudo-scalar Higgs are currently not supported in the official version of `aMCSusHi`. A user interested in decaying the pseudo-scalar Higgs is strongly encouraged to contact us.

The other parameters are relevant to `SusHi`. `Block sushi[1]` chooses the model with the three options SM (`Block sushi[1]`$= 0$), MSSM (`Block sushi[1]`$= 1$) and 2HDM (`Block sushi[1]`$= 2$). The second entry of `Block sushi` determines the Higgs boson: light Higgs (`Block sushi[2]`$=$ 11 or 0), pseudo-scalar Higgs (`Block sushi[2]`$=$ 21 or 1) and heavy Higgs (`Block sushi[2]`$=$ 12 or 2). The choice of the masses of the relevant Higgs bosons depends on the model. As stated before, `Block mass[25]` sets the Higgs mass in the SM. In the 2HDM, this entry corresponds to the mass of the light Higgs boson, while `Block mass[35]` and `Block mass[36]` specify the input for the heavy and the pseudo-scalar Higgs, respectively. All other 2HDM inputs are set in the information for `SusHi`. For reference, we give an example of the corresponding inputs for a heavy Higgs in the 2HDM below:[9]

```
[...]

#####################################
## INFORMATION FOR SUSHI
#####################################
Block sushi
  1   2          # model: 0 = SM, 1 = MSSM, 2 = 2HDM
  2   2          # 0 = light Higgs (h), 1 = pseudoscalar (A), 2 = heavy Higgs (H)
```

[7] Note that only in the on-shell scheme the automatic `MadGraph5_aMC@NLO` reweighting for μ_R is functionable.
[8] Bear in mind that this has to taken into account to identify the Higgs in the analysis of the showered events.
[9] A link to `2HDMC` [99] with the corresponding input convention is currently not supported.

```
Block renormbot # Renormalization of the bottom sector
  1   0         # m_b used for bottom Yukawa:  0 = OS, 1 = MSbar(m_b), 2 = MSbar(muR)
  2   2         # tan(beta)-res. of Y_b: 0 = no, 1 = naive, 2 = full (for OS only)
  4   4.75d0    # mbOS fixed
Block 2hdm # 2HDM version according to arXiv:1106.0034
  2             # (1=I,2=II ,3=III ,4=IV)
Block minpar
  3   50d0      # tanb
Block alpha
  0.0247d0      # mixing in Higgs sector
Block factors
  1   0d0       # factor for yukawa-couplings: c
  2   1d0       # t
  3   1d0       # b
```

Additionally to the inputs which we defined already for the SM the following parameters have to be set in the 2HDM: `Block renormbot[2]` specifies whether or not a resummation[10] of terms enhanced by $\tan(\beta)$ is applied through reweighting of the bottom Yukawa coupling as described in Ref. [74]; `Block 2hdm` determines which type of the 2HDM is used; the value of $\tan(\beta)$ is set through `Block minpar[3]`; and the mixing angle α corresponds to the entry in `Block alpha`.

The computation of MSSM Higgs cross sections requires `Block extpar`, `Block feynhiggs` and `Block renormsbot` in addition, which fix the parameters of the third family of quarks and squarks, determine the `FeynHiggs` inputs and yield information on the renormalization of the sbottom section, respectively. We will not provide any further information on these blocks, instead, we refer to the `SusHi` manual [106] and the `FeynHiggs` man pages [107]. Moreover, `Block alpha` can be omitted in the MSSM and the Higgs masses in `Block mass` have no effect, since they are determined by `FeynHiggs`, once `Block feynhiggs` is present. The MSSM Higgs mass that has been computed and applied in the run is provided to the user in `Block mass[25]` of the parameter card, which will be overwritten by the mass of the respective Higgs boson at the beginning of each MSSM run.

So far we did not comment on the `Block factors`. It allows to turn on and off individual contributions in all models. In fact, it even provides the possibility to rescale the respective Yukawa couplings by choosing values different from zero and one. With `Block factors[1]` one can include the charm quark in the computation. This requires to specify its $\overline{\text{MS}}$ mass $m_c(m_c)$ in `Block sminputs[8]` which is then translated to its on-shell mass. Furthermore, `Block factors[2]` and `Block factors[3]` multiply the top and the bottom Yukawa, respectively. In the MSSM, the stop Yukawa is rescaled by `Block factors[4]` and the sbottom one by `Block factors[5]`.

For further information on the input cards we refer the reader to Ref. [73] of `MadGraph5_aMC@NLO`, the manual of `SusHi` [106] and the man pages of `FeynHiggs` [107]. Three example parameter cards are provided in the folder <ggH-folder>/Cards; one for the SM (param_card.dat_SM), the 2HDM (param_card.dat_2HDM_scenB) and the MSSM (param_card.dat_MSSM_mhmodp). They match the scenarios that we study in the result section of this paper.

[10] See Refs. [100–105] for further information.

3.3 Choosing different shower scales

The choice of the shower scale is a very peculiar one in the gluon fusion process. In presence of the bottom-quark loop, factorization of soft and collinear radiation maybe spoiled at scales significantly smaller than the Higgs boson mass. This was pointed out by Ref. [69] in the context of analytic transverse momentum resummation. On the other hand, these terms might well be treated as a finite remainder as long as their impact remains moderate [68].

Due to their additive matching of the resummed low-p_T region with the fixed-order distribution valid at large transverse momenta, analytic p_T-resummation and the MC@NLO method are quite similar. In both cases there is a scale associated with that matching, the resummation scale $Q_{\rm res}$ and the shower scale $Q_{\rm sh}$, respectively. These scales can be interpreted as transition scales that separate the soft/collinear from the hard physics, very similar to the factorization scale of the PDFs. In other words, they define the range where resummation, and therefore the shower in MC@NLO, takes effect. Their value has to be chosen of the order of the typical scale of the problem.

In gluon fusion, the typical scale depends on the quark considered in the loop. Since $m_t \sim m_\phi$, there exist only two relevant scales for the top-quark loop (m_ϕ and p_T) and the shower scale can be chosen of the order of the Higgs mass. When considering the bottom loop, on the other hand, we face a three-scale problem (m_ϕ, m_b and p_T) which has not been solved to date. However, it has been suggested [69] to apply a lower transition scale to the bottom contribution, which, in particular, respects the fact that soft/collinear factorization is valid only up to smaller scales for the bottom loop. In Ref. [72] it was further proposed to separate three contributions according to their Yukawa couplings: the square of the top and the bottom, and their interference; and choose separate shower/resummation scales for all of them. This splitting allows for a model independent treatment of the problem by a rescaling of the individual contributions with the respective top and bottom Yukawas of a specific scenario in the 2HDM as well as the MSSM when neglecting squark effects. In the literature, two pragmatic approaches with physical motivation have been presented [72, 108] to determine separate scales for the three contributions. Their comparison will be studied elsewhere [109]. When studying phenomenological results in Section 4 we will apply the scales from Ref. [72] (referred to as "HMW" in what follows).

The separation of the bottom contribution (including the interference) from the top one with different shower scales (Q_b and Q_t, respectively) requires three runs in `MadGraph5_aMC@NLO`, which have to be combined as follows:

$$\sigma(Q_t, Q_b) = \sigma_t(Q_t) + \sigma_{t+b}(Q_b) - \sigma_t(Q_b). \tag{2}$$

To obtain all three contributions of different Yukawa origin with different scales that allows for a model independent treatment, on the other hand, `MadGraph5_aMC@NLO` has to be run five times:

$$\sigma(Q_t, Q_b, Q_{tb}) = \sigma_t(Q_t) + \sigma_b(Q_b) + \sigma_{t+b}(Q_{tb}) - \sigma_t(Q_{tb}) - \sigma_b(Q_{tb}). \tag{3}$$

The scales Q_t, Q_b and Q_{tb} determine the scale for top, the bottom and their interference,

respectively. As indicated before, the individual contributions can be separated using the `Block factors` in the parameter card.

The shower scale in `MadGraph5_aMC@NLO` cannot simply be accessed through the input cards, since it requires an advanced user to be familiar with its specific treatment in the code. `MadGraph5_aMC@NLO` does not use a simple fixed scale for Q_{sh}, instead, it statistically extracts the shower scale from a distribution peaked at a specific value. The user can only change the range of the interval of the distribution which of course also affects the peak. Therefore, we identify Q_t, Q_b and Q_{tb} in Eqs. (2) and (3) with the peak of the respective shower scale distributions.

The so-called shape parameters define the interval of the distribution from which the shower scale is picked on a event-wise basis. They can be specified in the include file <ggH-folder>/SubProcesses/madfks_mcatnlo.inc, where the relevant part is given by (`MadGraph5_aMC@NLO` default values):

```
[...]

c Define lower and upper veto range (see MC subtraction terms)
      double precision frac_low,frac_upp
      parameter (frac_low=0.1d0)
      parameter (frac_upp=1.0d0)

c Other control switches on veto (see MC subtraction terms)
      double precision scaleMClow,scaleMCdelta
      parameter (scaleMClow=10.d0)
      parameter (scaleMCdelta=20.d0)

[...]
```

The parameters `frac_low` and `frac_upp` are used to compute the upper and lower bounds of the Q_{sh} distribution, which will be explained in more detail below, while `scaleMClow` allows to set an absolute value of the lower bound on Q_{sh} and `scaleMCdelta` is used to apply a minimal value to the size of the distribution interval. In formulas the interval is defined by[11]

$$\begin{aligned} &Q_{\text{min}} \leq Q_{\text{sh}} \leq Q_{\text{max}}, \quad \text{with} \\ &Q_{\text{min}} = \max(\texttt{frac_low} \cdot \sqrt{s_0}, \texttt{scaleMClow}) \quad \text{and} \\ &Q_{\text{max}} = \max(\texttt{frac_upp} \cdot \sqrt{s_0}, Q_{\text{min}} + \texttt{scaleMCdelta}), \end{aligned} \tag{4}$$

where s_0 is the Born-level partonic center of mass energy squared. Evidently, `scaleMClow` and `scaleMCdelta` only take effect if the interval obtained through `frac_low` and `frac_upp` does not meet the corresponding restrictions. The corresponding Q_{sh} distribution is peaked around

$$Q_{\text{peak}} \sim \frac{(\texttt{frac_low} + \texttt{frac_upp})}{2} \sqrt{\langle s_0 \rangle}. \tag{5}$$

For a $2 \to 1$ process like gluon fusion this relation is an identity and $\sqrt{s_0}$ equals the mass of the final state particle, i. e., the Higgs mass in the case of gluon fusion. To change the peak-value to its half, e. g., for shower scale variations, one can simply divide `frac_low` and `frac_upp` by a

[11] For further information we refer the reader to Section 2.4.4 of Ref. [73].

factor of two. In this sense, it is convenient to keep the ratio between `frac_low` and `frac_upp` a constant, which in the default setup of `MadGraph5_aMC@NLO` is a factor of ten. Under this prerequisite, in order to choose a specific shower scale $Q_{\rm sh}$ for $gg \to \phi$ we simply have to determine

$$\texttt{frac_upp} = \frac{2}{1.1} \cdot \frac{Q_{\rm sh}}{m_\phi} \quad \text{and} \quad \texttt{frac_low} = \frac{\texttt{frac_upp}}{10}. \tag{6}$$

Here and in what follows, we associate $Q_{\rm peak}$ with the shower scale $Q_{\rm sh}$ and vice versa unless stated otherwise. After modifying the corresponding include file accordingly, `MadGraph5_aMC@NLO` has to be recompiled. This can be achieved by typing

```
> make clean
```

inside the <ggH-folder> which forces a recompilation during the next run of the code.

In Section 4 we show some applications of the `aMCSusHi` code and study the effect of different treatments of the shower scales.

4 Results: brief application

The $gg \to \phi$ process folder created by the `aMCSusHi` script preserves all the highly convenient features that come with `MadGraph5_aMC@NLO`. Besides many others, this entails an interface to the most common showers, the fully-automatic determination of scale and PDF variations without any extra-costs of computing time [110], the creation of any number of observables with a single run and analysis routines available for the most important processes including the gluon fusion Higgs production mode.

`aMCSusHi` allows to compute gluon-induced Higgs production including the complete dependence on the quark masses in the SM for the first time in a MC@NLO-type matching applying all versions of `Pythia` and `Herwig` Monte Carlos. While previous computations did only feature the `Herwig` showers [65–67], phenomenological results exist to our knowledge only for `Herwig6` [111]. As a first application we therefore study the impact of different showers on the top- and bottom-mass effects with respect to the heavy-top approximation at the 13 TeV LHC. For this purpose, Fig. 2 shows the ratio of the NLO+PS computation including mass effects and the corresponding cross section in the heavy-top limit as a function of the transverse momentum of the Higgs for different Monte Carlos: `Pythia8` (black, solid), `Herwig++` (red, dotted), `Pythia6` p_T-ordered (blue, dashed with points), `Pythia6` Q-ordered (green, dash-dotted with open boxes) and `Herwig6` (yellow, solid with filled boxes). We apply the MSTW2008 68%CL NLO PDF set [112] with the corresponding value of the strong coupling constant. The shower scale has been chosen as $Q_{\rm sh} = m_h/2$ in all cases, while for $\mu_{\rm F}$ and $\mu_{\rm R}$ we use the defaults specified in Section 3.2. Clearly, the mass effects are hardly dependent on the specific Monte Carlo which is particularly evident at small ($p_T \lesssim 50\,\text{GeV}$) and large ($p_T \gtrsim 150\,\text{GeV}$) transverse momenta. Nevertheless, there are some visible differences in the intermediate region which consistently discriminate the `Herwig` from the `Pythia` showers. Overall, they are at most 5% though and therefore still moderate.



Figure 2: Transverse momentum distribution of a SM Higgs at NLO+PS in the full theory normalized to the one in the heavy top effective field theory for different Monte Carlos: `Pythia8` (black solid), `Herwig++` (red, dotted), `Pythia6` p_T-ordered (blue, dashed with points), `Pythia6` Q-ordered (green, dash-dotted with open boxes) and `Herwig6` (yellow, solid with filled boxes).

Fig. 3 shows the effects of quark masses with respect to the heavy-top approximation as well, but for different choices of the associated shower scales. In all cases, the denominator and therefore the distribution in the heavy-top limit is computed with the respective scale of the top contribution $Q_{\rm sh} = Q_t$. As we observed before the Monte Carlo dependence is quite small; therefore, we only consider the `Pythia8` shower. For reference the black solid curve is the same as in Fig. 2 with $Q_{\rm sh} = m_h/2$ for all contributions. We compare it to the scales choices proposed in Ref. [69], which imply setting the shower scale of the bottom contribution (including the interference) to the bottom mass following Eq. (2) (red dotted curve). For the blue dashed curve with points we chose the HMW scales determined in Ref. [72] which can be found in Table 1 of that paper, applying a three-scale approach according to Eq. (3) ($Q_t = 49\,\text{GeV}$, $Q_{tb} = 34\,\text{GeV}$ and $Q_b = 23\,\text{GeV}$). The green dash-dotted curve with open boxes serves mostly for comparison with previous `Herwig6` results [111] which were computed with the scales of Ref. [69] as well.

For the p_T distribution in Fig. 3 (a), the change of the scale of the bottom contribution to $Q_b = m_b$ has a significant impact on the mass effects at small and intermediate transverse momenta. It



(a) (b)

Figure 3: (a) Same as Fig. 2, but for different choices of the shower scales, see text for details; (b) corresponding plot for the rapidity distribution of the Higgs.

develops an extremely steep drop at small transverse momenta which due to unitarity affects also the intermediate p_T-range in the opposite direction. The benefit of the usage of such a low scale is clearly disputable. While the `Herwig6` curve agrees rather well with previous result of Ref. [111] becoming flat for $p_T \lesssim 5\,\text{GeV}$, the `Pythia8` curve develops a steep increase in this region. This signals a significant Monte Carlo dependence at very small p_T which is not observed for larger Q_b scales. Furthermore, the rigorously low value also poses a technical problem in the code regarding the fact that the default shower scale choice in `MadGraph5_aMC@NLO`, as explained in Section 3.3, is a distribution. In order to solve this problem, we had to use a fixed value of $Q_{\text{sh}} = Q_b$ by setting `frac_low` = `frac_upp` = Q_b/m_h and `scaleMClow` = `scaleMCdelta` = 0.

Considering the HMW scales, the scale of the bottom contribution is not chosen at such low values. We find that the mass effects in this case (blue dashed line with points) are rather similar to the ones where all scales are set to $m_h/2$ (black solid line), although the individual HMW scales being quite different from this value. Looking at the rapidity distribution in Fig. 3 (b), on the other hand, we observe the expected feature of being essentially insensitive to any choice of the respective shower scales. We shall note at this point that simply due to their inclusion in the default analysis we were able to produce a large number of further observables at no additional computing cost.

To demonstrate the range of applicability of `aMCSusHi`, we consider two realistic BSM scenarios in Fig. 4 and Fig. 5: the heavy Higgs boson in Scenario B of Ref. [113] (a bottom dominated 2HDM



(a) (b)

Figure 4: Transverse momentum distribution of (a) the heavy Higgs boson and (b) the associated hardest jet computed in a bottom dominated scenario of the 2HDM (see text for details). Graphical notation is the following: black solid curve shows `Pythia8` at NLO+PS, red dotted curve is the same at LO+PS (normalized the NLO) and the blue dashed one with points corresponds to the fixed order curve at NLO.

scenario); and the pseudo-scalar Higgs boson in the $m_h^{\rm mod+}$(800,40) MSSM scenario [114] defined in Table 2 of Ref. [72]. The corresponding input files can be found in the folder <ggH-folder>/Cards.

In Fig. 4 we study the transverse momentum distributions of the Higgs and the hardest jet, while Fig. 5 depicts their rapidity distributions. In both cases we apply the HMW scales of Ref. [72]. At low transverse momenta the p_T distributions have similar shapes comparing the red dotted (LO+PS) to the black solid curves (NLO+PS). This can be easily inferred from the first inset where all curves are normalized to the black solid line in the main frame. However, it is well known that at the LO+PS p_T distributions yield unphysical results for transverse momenta beyond the shower scales indicated by a steep drop. Note that both curves are normalized to the same (the NLO) cross section. Continuing the comparison at hand, we observe a significant reduction of the scale uncertainties shown in the lower inset, where the bands are obtained by dividing the upper and lower bound of the respective cross section by the same central cross



(a) (b)

Figure 5: Rapidity distribution of (a) the pseudo-scalar Higgs boson and (b) the associated hardest jet computed in the $m_h^{\text{mod+}}$ scenario [114] of the MSSM with $M_A = 800\,\text{GeV}$ und $\tan\beta = 40$. Graphical notation is the same as in Fig. 4. All curves are normalized so that their bins add up to one.

section as in the first inset. The uncertainties correspond to the independent variation of all unphysical scales (μ_R, μ_F, Q_t, Q_b, Q_{tb}) by a factor of two. Comparing NLO+PS to the NLO fixed-order result denoted by fNLO, we observe the expected matching towards large transverse momenta.

In order to compare shapes, the rapidity distributions in Fig. 5 are normalized in a way that the sum of their bins yields one. We see that for the Higgs rapidity in Fig. 5 (a) all curves agree extremely well in terms of shape up to the forward region in which, nevertheless, the deviations are still well within the respective uncertainty bands.[12] For the rapidity distribution of the hardest jet the same is true for the two showered results, while the fNLO distribution, on the other hand, agrees only in the central region $|y(j_1)| \lesssim 3$, but features a significant enhancement when the hardest jet is more forward. In this region the cross section will receive large effects of collinear radiation which renders the shower to yield the more reliable description.

[12] Note that the Higgs reaches its kinematical limit slightly before $|y(\phi)| \sim 3$ already.

5 Conclusions

In this article we presented the new tool `aMCSusHi` which is a link between `MadGraph5_aMC@NLO` and `SusHi` for the computation of Higgs cross sections in gluon fusion at hadron colliders. The code gives NLO+PS accurate results in the SM, 2HDM and MSSM. The inputs in the MSSM are conveniently controlled through a link to `FeynHiggs`. We discussed the specific treatment of the shower scale in `MadGraph5_aMC@NLO` and pointed out its special role in the context of gluon-induced Higgs production. In the phenomenological part we study the impact of different shower scale choices on the mass effects in the SM. Furthermore, we studied results for 2HDM and MSSM Higgs production as an application of `aMCSusHi`. The `aMCSusHi` script can be downloaded from `https://cp3.irmp.ucl.ac.be/projects/madgraph/wiki/HSushi#no1`.

As an outlook, one may improve the SM prediction for the Higgs production mode through gluon fusion by merging the NLO+PS cross section for $gg \to h$ in the full theory with higher multiplicities computed in the heavy-top effective field theory. This is certainly beyond the scope of the present paper and is left for a future publication.

Acknowledgements. We are indebted to Emanuele Bagnaschi, Stefano Frixione, Robert Harlander, Paolo Torrielli and Alessandro Vicini for several fruitful discussions and comments on the manuscript. We would like to thank Marco Zaro and Rikkert Frederix for their help on technical issues concerning `MadGraph5_aMC@NLO` and Stefan Liebler for providing the diagrams in Fig. 1. HM was supported by a Marie Curie Early Initial Training Network Fellowship of the European Community's Seventh Framework Programme under contract number (PITN-GA-2012-315877-MCnetITN). This research was supported in part by the Swiss National Science Foundation (SNF) under contract 200020-141360.

References

[1] **ATLAS** Collaboration, G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, Phys.Lett. **B716** (2012) 1–29, [`arXiv:1207.7214`].

[2] **CMS** Collaboration, S. Chatrchyan et al., *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, Phys.Lett. **B716** (2012) 30–61, [`arXiv:1207.7235`].

[3] **LHC Higgs Cross Section Working Group** Collaboration, S. Dittmaier et al., *Handbook of LHC Higgs Cross Sections: 1. Inclusive Observables*, `arXiv:1101.0593`.

[4] **LHC Higgs Cross Section Working Group** Collaboration, S. Dittmaier et al., *Handbook of LHC Higgs Cross Sections: 2. Differential Distributions*, `arXiv:1201.3084`.

[5] **LHC Higgs Cross Section Working Group** Collaboration, S. Heinemeyer et al., *Handbook of LHC Higgs Cross Sections: 3. Higgs Properties*, arXiv:1307.1347.

[6] H. Georgi, S. Glashow, M. Machacek, and D. V. Nanopoulos, *Higgs Bosons from Two Gluon Annihilation in Proton Proton Collisions*, Phys.Rev.Lett. **40** (1978) 692.

[7] S. Dawson, *Radiative corrections to Higgs boson production*, Nucl.Phys. **B359** (1991) 283–300.

[8] A. Djouadi, M. Spira, and P. Zerwas, *Production of Higgs bosons in proton colliders: QCD corrections*, Phys.Lett. **B264** (1991) 440–446.

[9] J. M. Campbell, R. K. Ellis, F. Maltoni, and S. Willenbrock, *Higgs-Boson production in association with a single bottom quark*, Phys.Rev. **D67** (2003) 095002, [hep-ph/0204093].

[10] R. V. Harlander and W. B. Kilgore, *Higgs boson production in bottom quark fusion at next-to-next-to leading order*, Phys.Rev. **D68** (2003) 013001, [hep-ph/0304035].

[11] S. Dittmaier, M. Krämer, and M. Spira, *Higgs radiation off bottom quarks at the Tevatron and the CERN LHC*, Phys.Rev. **D70** (2004) 074010, [hep-ph/0309204].

[12] S. Dawson, C. Jackson, L. Reina, and D. Wackeroth, *Exclusive Higgs boson production with bottom quarks at hadron colliders*, Phys.Rev. **D69** (2004) 074027, [hep-ph/0311067].

[13] R. V. Harlander, K. J. Ozeren, and M. Wiesemann, *Higgs plus jet production in bottom quark annihilation at next-to-leading order*, Phys.Lett. **B693** (2010) 269–273, [arXiv:1007.5411].

[14] R. Harlander and M. Wiesemann, *Jet-veto in bottom-quark induced Higgs production at next-to-next-to-leading order*, JHEP **1204** (2012) 066, [arXiv:1111.2182].

[15] S. Bühler, F. Herzog, A. Lazopoulos, and R. Müller, *The fully differential hadronic production of a Higgs boson via bottom quark fusion at NNLO*, JHEP **1207** (2012) 115, [arXiv:1204.4415].

[16] R. V. Harlander, A. Tripathi, and M. Wiesemann, *Higgs production in bottom quark annihilation: Transverse momentum distribution at NNLO+NNLL*, Phys.Rev. **D90** (2014), no. 1 015017, [arXiv:1403.7196].

[17] M. Wiesemann, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, and P. Torrielli, *Higgs production in association with bottom quarks*, JHEP **1502** (2015) 132, [arXiv:1409.5301].

[18] M. Spira, A. Djouadi, D. Graudenz, and P. Zerwas, *Higgs boson production at the LHC*, Nucl.Phys. **B453** (1995) 17–82, [hep-ph/9504378].

[19] R. Harlander and P. Kant, *Higgs production and decay: Analytic results at next-to-leading order QCD*, JHEP **0512** (2005) 015, [hep-ph/0509189].

[20] R. V. Harlander and M. Steinhauser, *Hadronic Higgs production and decay in supersymmetry at next-to-leading order*, Phys.Lett. **B574** (2003) 258–268, [hep-ph/0307346].

[21] R. V. Harlander and M. Steinhauser, *Supersymmetric Higgs production in gluon fusion at next-to-leading order*, JHEP **0409** (2004) 066, [hep-ph/0409010].

[22] R. Harlander and M. Steinhauser, *Effects of SUSY QCD in hadronic Higgs production at next-to-next-to-leading order*, Phys.Rev. **D68** (2003) 111701, [hep-ph/0308210].

[23] R. V. Harlander and F. Hofmann, *Pseudo-scalar Higgs production at next-to-leading order SUSY-QCD*, JHEP **0603** (2006) 050, [hep-ph/0507041].

[24] M. Mühlleitner and M. Spira, *Higgs Boson Production via Gluon Fusion: Squark Loops at NLO QCD*, Nucl.Phys. **B790** (2008) 1–27, [hep-ph/0612254].

[25] U. Aglietti, R. Bonciani, G. Degrassi, and A. Vicini, *Analytic Results for Virtual QCD Corrections to Higgs Production and Decay*, JHEP **0701** (2007) 021, [hep-ph/0611266].

[26] R. Bonciani, G. Degrassi, and A. Vicini, *Scalar particle contribution to Higgs production via gluon fusion at NLO*, JHEP **0711** (2007) 095, [arXiv:0709.4227].

[27] G. Degrassi and P. Slavich, *On the NLO QCD corrections to Higgs production and decay in the MSSM*, Nucl.Phys. **B805** (2008) 267–286, [arXiv:0806.1495].

[28] C. Anastasiou, S. Beerli, and A. Daleo, *The Two-loop QCD amplitude $gg \to h, H$ in the Minimal Supersymmetric Standard Model*, Phys.Rev.Lett. **100** (2008) 241806, [arXiv:0803.3065].

[29] R. V. Harlander, F. Hofmann, and H. Mantler, *Supersymmetric Higgs production in gluon fusion*, JHEP **1102** (2011) 055, [arXiv:1012.3361].

[30] G. Degrassi and P. Slavich, *NLO QCD bottom corrections to Higgs boson production in the MSSM*, JHEP **1011** (2010) 044, [arXiv:1007.3465].

[31] M. Mühlleitner, H. Rzehak, and M. Spira, *SUSY-QCD Corrections to MSSM Higgs Boson Production via Gluon fusion*, PoS **RADCOR2009** (2010) 043, [arXiv:1001.3214].

[32] G. Degrassi, S. Di Vita, and P. Slavich, *NLO QCD corrections to pseudoscalar Higgs production in the MSSM*, JHEP **1108** (2011) 128, [arXiv:1107.0914].

[33] E. Bagnaschi, G. Degrassi, P. Slavich, and A. Vicini, *Higgs production via gluon fusion in the POWHEG approach in the SM and in the MSSM*, JHEP **1202** (2012) 088, [arXiv:1111.2854].

[34] G. Degrassi, S. Di Vita, and P. Slavich, *On the NLO QCD Corrections to the Production of the Heaviest Neutral Higgs Scalar in the MSSM*, Eur.Phys.J. **C72** (2012) 2032, [arXiv:1204.1016].

[35] R. V. Harlander and W. B. Kilgore, *Next-to-next-to-leading order Higgs production at hadron colliders*, Phys.Rev.Lett. **88** (2002) 201801, [hep-ph/0201206].

[36] C. Anastasiou and K. Melnikov, *Higgs boson production at hadron colliders in NNLO QCD*, Nucl.Phys. **B646** (2002) 220–256, [hep-ph/0207004].

[37] V. Ravindran, J. Smith, and W. L. van Neerven, *NNLO corrections to the total cross-section for Higgs boson production in hadron hadron collisions*, Nucl.Phys. **B665** (2003) 325–366, [hep-ph/0302135].

[38] C. Anastasiou, K. Melnikov, and F. Petriello, *Higgs boson production at hadron colliders: Differential cross sections through next-to-next-to-leading order*, Phys.Rev.Lett. **93** (2004) 262002, [hep-ph/0409088].

[39] S. Catani and M. Grazzini, *An NNLO subtraction formalism in hadron collisions and its application to Higgs boson production at the LHC*, Phys.Rev.Lett. **98** (2007) 222002, [hep-ph/0703012].

[40] M. Grazzini, *NNLO predictions for the Higgs boson signal in the $H \to WW \to l\nu l\nu$ and $H \to ZZ \to 4l$ decay channels*, JHEP **0802** (2008) 043, [arXiv:0801.3232].

[41] S. Catani, D. de Florian, M. Grazzini, and P. Nason, *Soft gluon resummation for Higgs boson production at hadron colliders*, JHEP **0307** (2003) 028, [hep-ph/0306211].

[42] A. Idilbi, X.-d. Ji, J.-P. Ma, and F. Yuan, *Threshold resummation for Higgs production in effective field theory*, Phys.Rev. **D73** (2006) 077501, [hep-ph/0509294].

[43] V. Ravindran, *Higher-order threshold effects to inclusive processes in QCD*, Nucl.Phys. **B752** (2006) 173–196, [hep-ph/0603041].

[44] V. Ahrens, T. Becher, M. Neubert, and L. L. Yang, *Renormalization-Group Improved Prediction for Higgs Production at Hadron Colliders*, Eur.Phys.J. **C62** (2009) 333–353, [arXiv:0809.4283].

[45] A. Djouadi and P. Gambino, *Leading electroweak correction to Higgs boson production at proton colliders*, Phys.Rev.Lett. **73** (1994) 2528–2531, [hep-ph/9406432].

[46] G. Degrassi and F. Maltoni, *Two-loop electroweak corrections to Higgs production at hadron colliders*, Phys.Lett. **B600** (2004) 255–260, [hep-ph/0407249].

[47] C. Anastasiou, R. Boughezal, and F. Petriello, *Mixed QCD-electroweak corrections to Higgs boson production in gluon fusion*, JHEP **0904** (2009) 003, [arXiv:0811.3458].

[48] U. Aglietti, R. Bonciani, G. Degrassi, and A. Vicini, *Two loop light fermion contribution to Higgs production and decays*, Phys.Lett. **B595** (2004) 432–441, [hep-ph/0404071].

[49] S. Actis, G. Passarino, C. Sturm, and S. Uccirati, *NLO Electroweak Corrections to Higgs Boson Production at Hadron Colliders*, Phys.Lett. **B670** (2008) 12–17, [arXiv:0809.1301].

[50] R. Bonciani, G. Degrassi, and A. Vicini, *On the Generalized Harmonic Polylogarithms of One Complex Variable*, Comput.Phys.Commun. **182** (2011) 1253–1264, [arXiv:1007.1891].

[51] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog, and B. Mistlberger, *Higgs boson gluonfusion production at threshold in N^3LO QCD*, Phys.Lett. **B737** (2014) 325–328, [arXiv:1403.4616].

[52] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog, and B. Mistlberger, *Higgs boson gluon-fusion production beyond threshold in N3LO QCD*, arXiv:1411.3584.

[53] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog, and B. Mistlberger, *Higgs boson gluon-fusion production in N3LO QCD*, arXiv:1503.06056.

[54] S. Marzani, R. D. Ball, V. Del Duca, S. Forte, and A. Vicini, *Higgs production via gluon-gluon fusion with finite top mass beyond next-to-leading order*, Nucl.Phys. **B800** (2008) 127–145, [arXiv:0801.2544].

[55] R. V. Harlander and K. J. Ozeren, *Top mass effects in Higgs production at next-to-next-to-leading order QCD: Virtual corrections*, Phys.Lett. **B679** (2009) 467–472, [arXiv:0907.2997].

[56] R. V. Harlander and K. J. Ozeren, *Finite top mass effects for hadronic Higgs production at next-to-next-to-leading order*, JHEP **0911** (2009) 088, [arXiv:0909.3420].

[57] R. V. Harlander, H. Mantler, S. Marzani, and K. J. Ozeren, *Higgs production in gluon fusion at next-to-next-to-leading order QCD for finite top mass*, Eur.Phys.J. **C66** (2010) 359–372, [arXiv:0912.2104].

[58] A. Pak, M. Rogal, and M. Steinhauser, *Finite top quark mass effects in NNLO Higgs boson production at LHC*, JHEP **1002** (2010) 025, [arXiv:0911.4662].

[59] A. Pak, M. Rogal, and M. Steinhauser, *Production of scalar and pseudo-scalar Higgs bosons to next-to-next-to-leading order at hadron colliders*, JHEP **1109** (2011) 088, [arXiv:1107.3391].

[60] R. V. Harlander, T. Neumann, K. J. Ozeren, and M. Wiesemann, *Top-mass effects in differential Higgs production through gluon fusion at order α_s^4*, JHEP **1208** (2012) 139, [arXiv:1206.0157].

[61] T. Neumann and M. Wiesemann, *Finite top-mass effects in gluon-induced Higgs production with a jet-veto at NNLO*, JHEP **1411** (2014) 150, [arXiv:1408.6836].

[62] P. Nason, *A New method for combining NLO QCD with shower Monte Carlo algorithms*, JHEP **0411** (2004) 040, [hep-ph/0409146].

[63] H. Mantler and M. Wiesemann, *Top- and bottom-mass effects in hadronic Higgs production at small transverse momenta through LO+NLL*, Eur.Phys.J. **C73** (2013), no. 6 2467, [arXiv:1210.8263].

[64] S. Frixione and B. R. Webber, *Matching NLO QCD computations and parton shower simulations*, JHEP **0206** (2002) 029, [hep-ph/0204244].

[65] S. Frixione, F. Stoeckli, P. Torrielli, B. R. Webber, and C. D. White, *MC@NLO 4.10*, http://www.hep.phy.cam.ac.uk/theory/webber/MCatNLO/.

[66] S. Frixione, F. Stoeckli, P. Torrielli, B. R. Webber, and C. D. White, *The MC@NLO 4.0 Event Generator*, arXiv:1010.0819.

[67] S. Frixione, F. Stoeckli, P. Torrielli, and B. R. Webber, *NLO QCD corrections in Herwig++ with MC@NLO*, JHEP **1101** (2011) 053, [arXiv:1010.0568].

[68] A. Banfi, P. F. Monni, and G. Zanderighi, *Quark masses in Higgs production with a jet veto*, JHEP **1401** (2014) 097, [arXiv:1308.4634].

[69] M. Grazzini and H. Sargsyan, *Heavy-quark mass effects in Higgs boson production at the LHC*, JHEP **1309** (2013) 129, [arXiv:1306.4581].

[70] K. Hamilton, P. Nason, E. Re, and G. Zanderighi, *NNLOPS simulation of Higgs boson production*, JHEP **1310** (2013) 222, [arXiv:1309.0017].

[71] K. Hamilton, P. Nason, and G. Zanderighi, *Finite quark-mass effects in the NNLOPS POWHEG+MiNLO Higgs generator*, arXiv:1501.04637.

[72] R. V. Harlander, H. Mantler, and M. Wiesemann, *Transverse momentum resummation for Higgs production via gluon fusion in the MSSM*, JHEP **1411** (2014) 116, [arXiv:1409.0531].

[73] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H.-S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, JHEP **1407** (2014) 079, [arXiv:1405.0301].

[74] R. V. Harlander, S. Liebler, and H. Mantler, *SusHi: A program for the calculation of Higgs production in gluon fusion and bottom-quark annihilation in the Standard Model and the MSSM*, Comput.Phys.Commun. **184** (2013) 1605–1617, [arXiv:1212.3249].

[75] S. Liebler, *Neutral Higgs production at proton colliders in the CP-conserving NMSSM*, arXiv:1502.07972.

[76] G. Corcella, I. Knowles, G. Marchesini, S. Moretti, K. Odagiri, P. Richardson, M. H. Seymour, and B. R. Webber, *HERWIG 6: An Event generator for hadron emission reactions with interfering gluons (including supersymmetric processes)*, JHEP **0101** (2001) 010, [hep-ph/0011363].

[77] G. Corcella, I. Knowles, G. Marchesini, S. Moretti, K. Odagiri, P. Richardson, M. H. Seymour, and B. R. Webber, *HERWIG 6.5 release note*, hep-ph/0210213.

[78] T. Sjostrand, S. Mrenna, and P. Z. Skands, *PYTHIA 6.4 Physics and Manual*, JHEP **0605** (2006) 026, [hep-ph/0603175].

[79] T. Sjostrand, S. Mrenna, and P. Z. Skands, *A Brief Introduction to PYTHIA 8.1*, Comput.Phys.Commun. **178** (2008) 852–867, [arXiv:0710.3820].

[80] M. Bahr, S. Gieseke, M. Gigg, D. Grellscheid, K. Hamilton, O. Latunde-Dada, S. Platzer, and P. Richardson, *Herwig++ Physics and Manual*, Eur.Phys.J. **C58** (2008) 639–707, [arXiv:0803.0883].

[81] J. Bellm, S. Gieseke, D. Grellscheid, A. Papaefstathiou, S. Platzer, P. Richardson, C. Rohr, and T. Schuh, *Herwig++ 2.7 Release Note*, arXiv:1310.6877.

[82] C. Degrande, C. Duhr, B. Fuks, D. Grellscheid, O. Mattelaer, and T. Reiter, *UFO - The Universal FeynRules Output*, Comput.Phys.Commun. **183** (2012) 1201–1214, [arXiv:1108.2040].

[83] N. D. Christensen and C. Duhr, *FeynRules - Feynman rules made easy*, Comput.Phys.Commun. **180** (2009) 1614–1641, [arXiv:0806.4194].

[84] N. D. Christensen, P. de Aquino, C. Degrande, C. Duhr, B. Fuks, M. Herquet, F. Maltoni, and S. Schumann, *A Comprehensive approach to new physics simulations*, Eur.Phys.J. **C71** (2011) 1541, [arXiv:0906.2474].

[85] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr, and B. Fuks, *FeynRules 2.0 - A complete toolbox for tree-level phenomenology*, Comput.Phys.Commun. **185** (2014) 2250–2300, [arXiv:1310.1921].

[86] S. Heinemeyer, W. Hollik, and G. Weiglein, *FeynHiggs: A Program for the calculation of the masses of the neutral CP even Higgs bosons in the MSSM*, Comput.Phys.Commun. **124** (2000) 76–89, [hep-ph/9812320].

[87] S. Heinemeyer, W. Hollik, and G. Weiglein, *The Masses of the neutral CP - even Higgs bosons in the MSSM: Accurate analysis at the two loop level*, Eur.Phys.J. **C9** (1999) 343–366, [hep-ph/9812472].

[88] G. Degrassi, S. Heinemeyer, W. Hollik, P. Slavich, and G. Weiglein, *Towards high precision predictions for the MSSM Higgs sector*, Eur.Phys.J. **C28** (2003) 133–143, [hep-ph/0212020].

[89] M. Frank, T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, *The Higgs Boson Masses and Mixings of the Complex MSSM in the Feynman-Diagrammatic Approach*, JHEP **0702** (2007) 047, [hep-ph/0611326].

[90] T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, *High-Precision Predictions for the Light CP -Even Higgs Boson Mass of the Minimal Supersymmetric Standard Model*, Phys.Rev.Lett. **112** (2014), no. 14 141801, [arXiv:1312.4937].

[91] G. Degrassi, P. Slavich, and F. Zwirner, *On the neutral Higgs boson masses in the MSSM for arbitrary stop mixing*, Nucl.Phys. **B611** (2001) 403–422, [hep-ph/0105096].

[92] A. Brignole, G. Degrassi, P. Slavich, and F. Zwirner, *On the $\mathcal{O}(\alpha_t^2)$ two loop corrections to the neutral Higgs boson masses in the MSSM*, Nucl.Phys. **B631** (2002) 195–218, [hep-ph/0112177].

[93] A. Brignole, G. Degrassi, P. Slavich, and F. Zwirner, *On the two loop sbottom corrections to the neutral Higgs boson masses in the MSSM*, Nucl.Phys. **B643** (2002) 79–92, [hep-ph/0206101].

[94] A. Dedes, G. Degrassi, and P. Slavich, *On the two loop Yukawa corrections to the MSSM Higgs boson masses at large tan beta*, Nucl.Phys. **B672** (2003) 144–162, [hep-ph/0305127].

[95] S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, *High-precision predictions for the MSSM Higgs sector at $\mathcal{O}(\alpha_b\alpha_s)$*, Eur.Phys.J. **C39** (2005) 465–481, [hep-ph/0411114].

[96] S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, *The Higgs sector of the complex MSSM at two-loop order: QCD contributions*, Phys.Lett. **B652** (2007) 300–309, [arXiv:0705.0746].

[97] P. Z. Skands et al., *SUSY Les Houches accord: Interfacing SUSY spectrum calculators, decay packages, and event generators*, JHEP **0407** (2004) 036, [hep-ph/0311123].

[98] E. Bagnaschi, R. Harlander, S. Liebler, H. Mantler, P. Slavich, and A. Vicini, *Towards precise predictions for Higgs-boson production in the MSSM*, JHEP **1406** (2014) 167, [arXiv:1404.0327].

[99] D. Eriksson, J. Rathsman, and O. Stal, *2HDMC: Two-Higgs-Doublet Model Calculator Physics and Manual*, Comput.Phys.Commun. **181** (2010) 189–205, [arXiv:0902.0851].

[100] M. Carena, D. Garcia, U. Nierste, and C. E. Wagner, *Effective Lagrangian for the $\bar{t}bH^+$ interaction in the MSSM and charged Higgs phenomenology*, Nucl.Phys. **B577** (2000) 88–120, [hep-ph/9912516].

[101] J. Guasch, P. Hafliger, and M. Spira, *MSSM Higgs decays to bottom quark pairs revisited*, Phys.Rev. **D68** (2003) 115001, [hep-ph/0305101].

[102] D. Noth and M. Spira, *Supersymmetric Higgs Yukawa Couplings to Bottom Quarks at next-to-next-to-leading Order*, JHEP **1106** (2011) 084, [arXiv:1001.1935].

[103] D. Noth and M. Spira, *Higgs Boson Couplings to Bottom Quarks: Two-Loop Supersymmetry-QCD Corrections*, Phys.Rev.Lett. **101** (2008) 181801, [arXiv:0808.0087].

[104] L. Hofer, U. Nierste, and D. Scherer, *Resummation of tan-beta-enhanced supersymmetric loop corrections beyond the decoupling limit*, JHEP **0910** (2009) 081, [arXiv:0907.5408].

[105] L. Mihaila and C. Reisser, $\mathcal{O}(\alpha_s^2)$ *corrections to fermionic Higgs decays in the MSSM*, JHEP **1008** (2010) 021, [arXiv:1007.0693].

[106] R. V. Harlander, S. Liebler, and H. Mantler, *SusHi 1.5.0: A program for the calculation of Higgs production in gluon fusion and bottom-quark annihilation in the Standard Model, the 2HDM, the MSSM and the NMSSM*, http://sushi.hepforge.org/manual.html.

[107] *FeynHiggs man pages*, http://wwwth.mpp.mpg.de/members/heinemey/feynhiggs/cFeynHiggs.html.

[108] E. Bagnaschi and A. Vicini, *The Higgs transverse momentum distribution in gluon fusion as a multiscale problem*, to be published elsewhere.

[109] E. Bagnaschi, R. V. Harlander, H. Mantler, A. Vicini, and M. Wiesemann, *Resummation effects in Higgs transverse momentum distribution in the SM and beyond*, to be published elsewhere.

[110] R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, R. Pittau, and P. Torrielli, *Four-lepton production at hadron colliders: aMC@NLO predictions with theoretical uncertainties*, JHEP **1202** (2012) 099, [arXiv:1110.4738].

[111] S. Frixione, *talk given at the ggF meeting on Higgs pT*, CERN, July 2013, http://indico.cern.ch/event/263472/contribution/3/material/slides/.

[112] A. Martin, W. Stirling, R. Thorne, and G. Watt, *Parton distributions for the LHC*, Eur.Phys.J. **C63** (2009) 189–285, [arXiv:0901.0002].

[113] R. Harlander, M. Mühlleitner, J. Rathsman, M. Spira, and O. Stål, *Interim recommendations for the evaluation of Higgs production cross sections and branching ratios at the LHC in the Two-Higgs-Doublet Model*, arXiv:1312.5571.

[114] M. Carena, S. Heinemeyer, O. Stål, C. Wagner, and G. Weiglein, *MSSM Higgs Boson Searches at the LHC: Benchmark Scenarios after the Discovery of a Higgs-like Particle*, Eur.Phys.J. **C73** (2013), no. 9 2552, [arXiv:1302.7033].